## REMARKS

## I. Status of the Claims

Claims 1-8 are withdrawn.

Claims 10, 11, 13 and 14 are amended.

Claims 16 and 17 are new.

Claims 9-17 are pending.

## II. Summary

Applicant thanks Supervisor Padmanabhan and Examiner Cotton for suggestions on claim amendments to move this case toward allowance. A question of total daily dosage as distinct from unit dose, e.g. dose per capsule, arose. Claim amendments with support from the specification were suggested, as long as the claim scope was not within the art, e.g. in Examples 1-46 in Caruso. Support for the daily doses and unit doses are found on at least the following locations. Citations are to paragraphs in the specification:

## What is a "low dose" tricyclic antidepressant?

[00008] 25 mg/day or less.

[00010] 0.5 gm-2.6 gm daily; 0.5-2 gm/day acetaminophen; 0.6-2.6 gm/day aspirin; 0.6-1.8 gm/day ibuprofen.

## What is the "standard dose" of non-narcotic analgesic?

[00011] 2.5 mg to 25 mg/day (.5 mg to 2 mg, 10-15 mg/day).

[00015] Example 1: 500 mg acetaminophen + 5 mg doxepin (unit dose).

[00016] Example 2: 5 mg doxepin + 650 mg aspirin, unit dose, twice daily.

[00017] Example 3: 10 mg doxepin + 600 mg ibuprofen.

The references requested by the examiner regarding low and standard doses are in Exhibits A and B.

No fees are believed due at this time, however, please charge any additional deficiencies or credit any overpayments to deposit account number 12-0913 with reference to our attorney docket number (41957/102748).

Respectfully submitted,

Associated by Mosla (

Alice O. Martin Registration No. 35,601

April 12, 2007 Barnes & Thornburg LLP P.O. Box 2786 Chicago, IL 60690-2786

CHDS01 AOM 391366v1

## Harrison's

## PRINCIPLES OF INTERNAL MEDICINE

Fauci Braunwald Isselbacher Wilson Martin Kasper Hauser Longo Note: Dr. Fauci and Dr. Longo's works as editors and authors were performed outside the scope of their employment as U.S. government employees. These works represent their personal and professional views and not necessarily those of the U.S. government.

Harrison's
PRINCIPLES OF INTERNAL MEDICINE
Fourteenth Edition

Copyright © 1998, 1994, 1991, 1987, 1983, 1980, 1977, 1974, 1970, 1966, 1962, 1958 by The McGraw-Hill Companies, Inc. All rights reserved. Printed in the United States of America. Except as permitted under the United States Copyright Act of 1976, no part of this publication may be reproduced or distributed in any form or by any means, or stored in a data base or retrieval system, without the prior written permission of the publisher.

4567890 DOWDOW 0403020100 ISBN 0-07-020291-5 (COMBO) 0-07-020292-3 (VOL 1) 0-07-020293-1 (VOL 2) 0-07-912013-X (SET)

FOREIGN LANGUAGE EDITIONS
Arabic (Thireenth Edition)—McGraw-Hill Libri Italia sri (est. 1996)
Chinese (Twelfith Edition)—McGraw-Hill Bock Company-Singapore © 1994
Croatian (Thirteenth Edition)—McGraw-Hill Libri Italia sri (est. 1996)
French (Thirteenth Edition)—McGraw-Hill Libri Italia sri © 1995
German (Thirteenth Edition)—McGraw-Hill Libri Italia sri © 1995
German (Thirteenth Edition)—McGraw-Hill Libri Italia sri © 1995
Italian (Thirteenth Edition)—McGraw-Hill Libri Italia sri © 1995
Japanese (Eleventh Edition)—McGraw-Hill Libri Italia sri © 1995
Portuguese (Thirteenth Edition)—McGraw-Hill McGraw-Hill Interamericana © 1995
Spanish (Thirteenth Edition)—McGraw-Hill Interamericana © 1995
Spanish (Thirteenth Edition)—McGraw-Hill Interamericana © 1994

Turkish (Thirteenth Edition)-McGraw-Hill Libri Italia srl (est. 1996)

This book was set in Times Roman by Monotype Composition Company. The editors were I. Dereck Jeffers, Martin J. Wonsiewicz, and Mariapez Ramos Englis; editorial assistants were Ross Derario and Daniel J. Green. The production director was Robert Laffler. The index was prepared by Irving Tullar. The text and cover designer was Marsha CoherParallelogram Grashics.

R.R. Donnelley and Sons, Inc., was the printer and binder.

Library of Congress Cataloging-in-Publication Data

Harrison's principles of internal medicine.—14th. ed./editors, Anthony S. Fusci... [et al.] p. cm. Includes bibliographical references and index. ISBN 047-020291-5 (1 vol. cd.)—ISBN 047-912013-X (2 vol. ed. :sei)—ISBN 047-020292-3 (2 vol. ed. :bk 1)—ISBN 047-020293-1 (2 vol. ed. :bk 2).

U-07-U20293-1 (2 vol. ed. : bk 2)
 Internal medicine. 1. Harrison, Tinsley Randolph.
 II. Fauci, Anthony S.

[DNLM: 1. Internal Medicine. WB 115 H322 1998] RC46.H333 1998 616—dc21

DNLM/DLC for Library of Congress Dose, mg

650 PO

650 PO

The introduction of a parenteral form of NSAID, ketorolac, established sustibless of this class of compounds in the management of acute severe pain. Ketorolac is sufficiently potent and rapid in onset to supplant opioids for many patients with acute severe headache and musculoskeleal pain.

Opioid Analgesies: Opioids are the most potent pain-relieving drugs currently available. Furthermore, of all malagisacis, they have the broadest range of effices, providing the most reliable method for respiratory beginn, Although side effects are common, except for respiratory depression, they are usually not serious and can be reversed rapidly with the nacroic antagonist nalozone. The physician should not healtaste to use opioid analgesics in patients with acute sewere pain. Table 12-1 lists the most commond used notice datasets.

Opioids produce analgesia by actions in the central nervous system. They activate poin-inhibitory neurons and directly inhibit pain-transmission neurons. Most of the commercially available opioid analgesics act at the same opiate receptor (mu receptor), differing mainly in potency, speed of onset, duration of action, and optimal route of administration. Although the dose-related side effects (sedation, respiratory depression, purtues, constipation) are similar among the different opioids, some side effects are due to accumulation of ronopioid metabolites that are unique to individual drugs. One striking example of this is normeperidine, a metabolite of megeridine. Normeperidine produces hypersychalbility and seizures that are not

NONNARCOTIC ANALGESICS: USUAL DOSES AND INTERVALS

reversible with naloxone. Normeperidine accumulation is

The most rapid relief with opioids is obtained by introadministration; relief with oral administration is significantly as
Common acute side effects include nausea, vomiting, and sell
Those effects are dose-related, and there is great variability est
patients in the doses that relieve pain and produce side effects
and interval. The most important principle is to provide adequate,
relief. This requires asking the patient whether the drug has relie
the pain and, if so, when the relief wears off. The most cost,
error made by physicians in managing severe pain with opials
to prescribe an inadequate dose. Since many patients are reluct
to complain, this practice leaded to needless suffering. In the sixe
of sedation at the expected time of peak effect, a physician should
hesitate to repeat the initial dose to oschieve satisfactory pain 76
hesitate to repeat the initial dose to oschieve satisfactory pain 76
hesitate to repeat the initial dose to oschieve satisfactory pain 76

An innovative approach to the problem of achieving adequal neitled is the use of patient-controlled analgesis (PCA). It requires a device that immediately delivers a pre-programmed of an opioid drug when the patient pushes a button. The device be programmed to limit the total hourly dose so that overdosin impossible. The patient can then tituate the dose to the optimal be This approach is used most extensively for the management postoperative pain, but there is no reason why it should not be a for any hospitalized patient with persistent severe pain. PCI also used for home care of patients with intractable pain, such metastatic canner.

Enteric-coated preparations available

Side effects uncommon

Table 12-1 Drugs for Relief of Pain

Generic Name

Acetylsalicylic acid

Acetaminophen

Ibuprofen Naproxen Fenoprofen Indomethacin Ketorolac NARCOTIC ANALGESICS: USUA	400 PO 250–500 PO 200 PO 25–50 PO 15–60 IM	q 4–6 h q 12 h q 4–6 h q 8 h q 4–6 h	Available withour prescription Delayed effects may be due to long half-life pa Gastrointestinal side effects common Available for parenteral use (IM)	
Generic Name	Parenteral Dose, mg	PO Dose, mg	Comments	-
Codeine	30-60 q 4 h	30-60 q 4 h	Nausea common	5
Oxycodone	= •	5-10 q 4-6 h	Usually available with acetaminophen or aspirin	5
Morphine	10 q 4 h	60 q 4 <sup>°</sup> h		2
Morphine sustained release	pa ·	60-180 bid to tid	Oral slow-release preparation	Ē
Hydromorphone	1-2 q 4 h	2-4 q 4 h	Shorter acting than morphine sulfate	ጉ
Levorphanol	2 q 6-8 h	4 q 6-8 h	Longer acting than morphine sulfate; absorbed well Po	ñ
Methadone	10 q 6-8 h	20 q 68 h	Delayed sedation due to long half-life	×
Meperidine	75-100 q 3-4 h	300 q 4 h	Poorly absorbed PO; normeperidine a toxic metabolite	L
Butorphanol		1-2 q 4 h	Intranasal spray	,,
Fentanyl			Transdermal patch	c

Interval

a 4 h

q4h

### ANTICONVULSANTS AND ANTIARRHYTHMICS

Generic Name	PO Dose, mg	Interval
Phenytoin	300	daily/qhs
Carbamazepine	200-300	q 6 h
Clonazepam	1	q6h
Meriletine	150-300	a 6-12 h

## TRICYCLIC ANTIDEPRESSANTS

	Uptake	Blockade	0.1	Anticholinergic	Orthostatic	Cardiac	Average Dose,	Range.
Generic Name	5HT	NE	Sedative Potency	Potency	Hypotension	Arrhythmia	mg/day	mg/day
Doxepin	++	+	High	Moderate	Moderate	Less	200	75-400
Amitriptyline	++++	++	High	Highest	Moderate	Yes	150	25-30C
Imipramine	++++	++	Moderate	Moderate	High	Yes	200	75-40C
Nortriptyline	+++	++	Moderate	Moderate	Low	Yes	100	40-15C
Desipramine	+++	++++	Low	Low	Low	Yes	150	50-30C

# PHARMACOLOGICAL BASIS OF THERAPEUTICS

Joel G. Hardman Lee E. Limbird

Perry B. Molinoff Raymond W. Ruddon

Alfred Goodman Gilman

## Goodman and Gilman's THE PHARMACOLOGICAL BASIS OF THERAPEUTICS, 9/e

Copyright © 1996, 1990, 1985, 1980, 1975, 1970, 1965, 1955, 1941 by The McGraw-Hill Companies, Inc. All rights reserved. Primed in the United States of America. Except as perminded under the United States Copyright Act of 1976, no part of this publication may be reproduced or distributed in any form or by any means, or stored in a data base or retrieval system, without the prior written permission of the publisher.

234567890 DOWDOW 9987

ISBN 0-07-026266-7

This book was set in Times Roman by York Graphic Services, Inc. The editors were Martin J. Wonsiewicz and Peter McCurdy; the production supervisors were Robert Laffler and Clare Stanley, and the cover designer was Marsha Cohen/Paralellogram. The index was prepared by Irving Condé

R.R. Donnelley and Sons Company was printer and binder.

This book is printed on acid-free paper.

Library of Congress Cataloging-in-Publication Data

Goodman & Gilman's The Pharmacological Basis of Therapeutics. —9th ed. / Joel G. Hardman, Alfred Goodman Gilman, Lee E. Limbird.

p. cm.
 Includes bibliographical references and index.

Includes bibliographical references and index. ISBN 0-07-026266-7 (hardcover)

Pharmacology 2. Chemotherapy. I. Goodman, Louis Sanford. II. Gilman, Alfred.
 III. Hardman, Joel G. IV. Gilman, Alfred Goodman. V. Limbird, Lee E.

[DNLM: I. Pharmacology. 2. Drug Therapy. QV 4 G6532 1995] RM300.G644 1995

615' .7—dc20 DNLM/DLC

for Library of Congress

95-36658

there there is not the state of	pres-	pine) si pine	imip (inipolitic) olite) 1915 there 1956 2 tyline 25 8	pyclic lly to	ng the s and ients mod-	s iso. See the of the o	zines, chotic dents; accu.	com-c dative alogs, zines	benz- ily by even- chotic	in the It. yl de s, and
NONPROPRIETARY NAME (TRADE NAME)	DOSB AN	DOSE AND DOSAGE FORMS	RMS			aons	SLOB BPPECTS			
Norepiuephrlue-Reuptake Iuliibitors: Tertiary Amine Tricyclics										
	Usual	Extreme		, giri		Anni-				
R <sub>1</sub> R <sub>2</sub> R <sub>3</sub> · · · ·	Dose, mg/day	Dose, mg/day	Dosage Form	Amine Effects	Sedation	cholinergic Effects	Hypo- tension	Cardiac Effects	Seizures	Weight Gain
Amitriptyline (ELAVIL and others) C H C=CH(CH2)2N(CH3)2	100-200	25–300	0,1	NE, S-HT	+ + +	‡	‡ ‡	+ + +	<b>+</b>	‡
Clomigramine (Anafrana.) C Cl N—(CH2)3N(CH3)2	100-200	25-250	0	NE, S-HT	‡	‡ ‡	‡	+ + +	‡ ‡	+
Doxepin (ADAPIN, SINEQUAN) O H N=CH(CH <sub>2</sub> ) <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>	100-200	25-300	0	NE, S-HT	‡ ‡	‡	† †	+ +	+ +	‡
Impramine (TOFFANT, and others)  C H N—(CH <sub>2</sub> ) <sub>3</sub> N(CH <sub>3</sub> ) <sub>2</sub>	100-200	25-300	o, I	NE, S-HT	‡	<b>+</b>	‡	+ + +	<b>‡</b>	,‡
(+)-Trimipramine (surmontil.)  CH <sub>3</sub>										
C H N-CH2CHCH2N(CH3)2	75-200	25-300	0	NE, 5-HT	† †	+++	+	+++	‡	<b>+</b>
Norspinsplirius-Reuptake Inhibliors: Secondary Amine Tricyclics Amoxapine (ASENDR)	200-300	30-600	۰	NB, DA	+	+	‡	‡	‡	+
ÇZ"										
Designantine (NORPRANIK, PERTOFRANE)	100-200	25–300	•	뿐	0/+	+	+	· ‡	+	+
CH <sub>2</sub> CH <sub>2</sub> NHCH <sub>3</sub>										
Маргоцііпе (слоюми.)	100-150	25-225	0	N E	‡	‡	‡	÷	+ + +	+
CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> NHCH <sub>3</sub>										
								4		